

### **MEETING ABSTRACT**

**Open Access** 

# Tobacco smoking in eating disordered female patients

Rénate Eiber<sup>1,2</sup>

From 1<sup>st</sup> International Congress on Neurobiology and Clinical Psychopharmacology and European Psychiatric Association Conference on Treatment Guidance Thessaloniki, Greece. 19-22 November 2009

#### **Background**

Some smokers use tobacco in order to control weight. It is conceivable that eating disordered women smoke for that reason. Aim: to describe smoking features in eating disordered women; to show differences according to types of eating disorder and to identify factors explaining tobacco use in this population.

#### Materials and methods

A retrospective study including two hundred women diagnosed for eating disorders according to ICD-10 criteria. We gathered socio-demographic features and evaluated tobacco dependence with the Fagerström questionnaire (FTND).

#### Results

The body mass index was the same in all smoking and no smoking eating disordered patients. FTND score and daily consumption of cigarettes in eating disordered smokers showed significant differences according to diagnostic groups of eating disorders. Age of smoking the first cigarette, age of beginning daily tobacco smoking and reasons of smoking did not display any significant differences between diagnostic groups. Daily smoking was much more frequent in bulimics.

#### **Conclusions**

Our study did not allow to reveal that tobacco is used to control weight in eating disordered women.

#### **Author details**

<sup>1</sup>Elber Renate psychiatrist private exercice 16 chemin du Calquet 31100 Toulouse, France. <sup>2</sup>BERLIN Ivan department of pharmacology Pitié-Salpêtrière Hospital Paris, France.

<sup>1</sup>Elber Renate psychiatrist private exercice 16 chemin du Calquet 31100 Toulouse, France



Published: 22 April 2010

#### References

- Neumark-Sztainer D, Story M, Dixon LB, et al: Adolescents engaging in unhealthy weight control behaviors: are they at risk for other healthcompromising behaviors? Am J Public Health 1998, 88:952-5.
- Wiseman CV, Turco RM, Sunday SR, et al: Smoking and body image concerns in adolescent girls. Int J Eat Disord 1998, 24:429-33.
- Stock SL, Goldberg E, Corbett S, et al: Substance use in female adolescents with eating disorders. J Adolesc Health 2002, 31:176-82.
- Anzengruber D, Klump KL, Thornton L, et al: Smoking in eating disorders. Fating Behaviors 2006. 7:291-99.

doi:10.1186/1744-859X-9-S1-S87

Cite this article as: Eiber: Tobacco smoking in eating disordered female patients. *Annals of General Psychiatry* 2010 **9**(Suppl 1):587.

## Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at www.biomedcentral.com/submit

